





BOGOLIUBOV LABORATORY OF THEORETICAL PHYSICS

History and researches





**The agreement on the establishment of
JINR
was signed on 26 March 1956 in Moscow**



Laboratory of Theoretical Physics, JINR

2 Decrees, May 25, 1956

П Р И К А З

ПО ЛИЧНОМУ СОСТАВУ ОБЪЕДИНЕННОГО ИНСТИТУТА

№ 6

"25" мая 1956 г.

- ЗАЧИСЛИТЬ:
1. БОГОЛЮБОВА Николая Николаевича временно начальником сектора № 3 Теоретической лаборатории с окладом 6000 руб. в месяц, с 1 июня с.г.
 2. ШИРКОВА Дмитрия Васильевича старшим научным сотрудником сектора № 3 Теоретической лаборатории с окладом 1500 руб. в месяц по совместительству, с 1 июня с.г.
 3. МЕДВЕДЕВА Бориса Валентиновича старшим научным сотрудником сектора № 3 Теоретической лаборатории с окладом 1500 руб. в месяц по совместительству, с 1 июня с.г.
 4. ПОЛИВАНОВА Михаила Константиновича научным сотрудником сектора № 3 Теоретической лаборатории с окладом 1000 руб. в месяц по совместительству, с 1 июня с.г.

ДИРЕКТОР

ОБЪЕДИНЕННОГО ИНСТИТУТА ЯДЕРНЫХ ИССЛЕДОВАНИЙ

Д.И. БЛОХИНЦЕВ
Д.И. БЛОХИНЦЕВ

П Р И К А З

ПО ЛИЧНОМУ СОСТАВУ ОБЪЕДИНЕННОГО ИНСТИТУТА

№ 5

"25" мая 1956 года.

До утверждения новой структуры Института возложить на академика БОГОЛЮБОВА Николая Николаевича /начальника сектора № 3 Теоретической лаборатории/ исполнение обязанностей директора Теоретической лаборатории Объединенного Института.

ДИРЕКТОР

ОБЪЕДИНЕННОГО ИНСТИТУТА ЯДЕРНЫХ ИССЛЕДОВАНИЙ

Д.И. БЛОХИНЦЕВ
Д.И. БЛОХИНЦЕВ


Nikolai Nikolaevich Bogoliubov (1909 – 1992)


is a distinguished scientist in the field of physics and mathematics. His scientific activity began in Kiev (1923-1947) And then continued in Moscow (since 1949) and Dubna (since 1956). Main scientific results are:

- Nonlinear mechanics: asymptotic methods, stability theory ;
- Statistical physics: kinetic equations, quasiaverages for systems with spontaneously broken symmetries;
- Quantum statistics: microscopic theory of Bose-gas superfluidity, microscopic theory of superconductivity ;
- Quantum field theory: axiomatic scattering matrix, general renormalization theory, renormalization group theory, proof of dispersion relations;
- Elementary Particle Theory: "quark bag" model, quantum number "colour".

N.N. Bogoliubov's scientific activity began at the age of 14 –15. His major independent results were obtained when he was 20–25.

N.N. Bogoliubov's scientific activity is specified by considerable mathematical culture and directness to solution of concrete problems of natural science.





Dmitrii Ivanovich Blokhintsev

(11.01. 1908 – 27.01.1979),

one of the pioneers of atomic science and technology in USSR, the organizer and the first director of the JINR.

Main scientific results in the fields:

- Quantum mechanics
- Acoustics of an inhomogeneous moving medium
- Neutron physics
- Quantum field theory
- Particle physics


1954 – Scientific supervisor of creation and putting into operation of the world first atomic power station

1956 – 1965 – JINR Director


1965–1979 – Director of Lab of Theoretical Physics




M.A. Markov (EPhL AN)



N.N. Bogoliubov



D.I. Blokhintsev




Ya.A. Smorodinsky (INP AN)

N.N. Bogoliubov
and his disciples







D.V. Shirkov



I.T. Todorov



V.G. Kadyshevsky



Nguyen Van Hieu



V.A. Matveev




A.N. Tavkhelidze



A.A. Logunov



V.A.Ambartsumian and N.N.Bogoliubov (1979)




**BLTP:
Fundamental Science
International Cooperation
and Education**



Scientific Policy:


Multidisciplinary Researches, Theoretical Physics on the Basis of Advanced Mathematics, Theoretical Support of the JINR Experimental Programs, Interplay of Scientific Researches and Education



Scientific Personnel

	D	C	O	Total
Fundamental Interactions	30	27	26	83
Nuclear Physics	25	23	20	68
Condensed Matter	16	22	10	48
Mathematical Physics	13	12	7	32
Total	84	84	63	231

Age structure



2018

SCIENTIFIC PERSONNEL BY COUNTRY (BLTP)

Country	Total	Country	Total
Russia	165	Germany	6
Czech Republic	5	Poland	3
Mongolia	1	Bulgaria	7
Turkey	1	Korea	1
Belarus	2	India	3
Kazakhstan	9	Uzbekistan	3
Slovakia	7	Moldavia	1
Azerbaijan	2	Mexico	1
Ukraine	3	Romania	3
Tajikistan	1	Japan	1
Vietnam	1	Argentina	1
Armenia	4		
Total - 231 (165 scientists from Russia and 66 from other countries)			

INTERNATIONAL SCIENTIFIC PERSONNEL

BY COUNTRY AND TOPIC

More than 20 countries are represented
More than 25% are from other than RF member states

Country	Fundamental Interactions	Nuclear Theory	Condensed Matter	Modern Mathematical Physics	Total
Total (RF/Other)	80 (54/26)	69 (51/18)	49 (37/12)	31 (23/8)	229 (165/64)

Web of Science®

JINR publication statistics

2011 – 2017

Total number of publications: 8 178
Total number of citations: 97 711
Excluding self-citations: 77 861
Average citations per article: 11,95
h-index: 106

2017

Total number of publications: 1 260
Total number of citations: 1 202
Excluding self-citations: 837
Average citations per article: 0,95
h-index: 14

2017: JINR in comparison with CERN

JINR

Total number of publications: 1 260
Total number of citations: 1 202
Excluding self-citations: 837
Average citations per article: 0,95
h-index: 14

CERN

Total number of publications: 1 287
Total number of citations: 1 694
Excluding self-citations: 1 226
Average citations per article: 1,32
h-index: 15

BLTP PUBLICATIONS (2013-2017)

	2013	2014	2015	2016	2017	Total
Journal publications	363	364	356	370	390	1843
Conference proceedings	122	131	142	165	170	730
Total	485	495	498	535	560	2573

2019-2023: Themes and projects

Theory of Fundamental Interactions

Projects:

- Standard Model and Its Extension,
- QCD Parton Distributions for Modern and Future Colliders,
- Physics of Heavy and Exotic Hadrons,
- Hadronic matter under extreme conditions

Theory of Condensed Matter

Projects:

- Physical properties of complex materials and nanostructures
- Mathematical problems of many-particle systems

Research and Education Project “Dubna International School of Theoretical Physics (DIAS-TH)”

Theory of Nuclear Structure and Nuclear Reactions

Projects:

- Nuclear Structure far from Stability Valley
- Nucleus-Nucleus Collisions and Nuclear Properties
- Exotic Few-Body Systems,
- Nuclear Structure and Dynamics at the Relativistic Energies.

Modern Mathematical Physics: Strings and Gravity, Supersymmetry, Integrability

Projects:

- Quantum groups and integrable systems
- Supersymmetry
- Quantum gravity, cosmology and strings

Theory of Fundamental Interactions

Leaders: D.I. Kazakov, O.V. Teryaev, A.B. Arbuzov

Further development of the **quantum field theory** approach in the framework of the Standard Model of fundamental interactions and its extensions. **Lattice simulations** for obtaining nonperturbative results in gauge theories. Elaboration of **multiloop calculations** in **QCD, Electroweak theory, and Minimal Supersymmetric Standard Model**. Theoretical predictions concerning the experimental observation of **supersymmetry**, the **Higgs boson**, investigation of the **spin structure of the nucleon**, **heavy flavor physics**, **vacuum structure in QCD**, and **hadron properties in dense and hot media**. Elaboration of new **phenomenological models to describe the hadron dynamics** in the framework of general principles of quantum field theory incorporating basic experimental patterns. **Theoretical support** of current and future **experiments at JINR, CERN, GSI, DESY, IHEP and other physics centers**.

Theory of Nuclear Structure and Nuclear Reactions

Leaders: V.V. Voronov, A.I. Vdovin, N.V. Antonenko

The main goals are to predict and analyze **properties of atomic nuclei beyond the stability valley** and investigate structure peculiarities of **superheavy and exotic nuclei**; to study dynamics of **nuclear collisions at low and intermediate energies** accompanied by production of stable as well as radioactive nuclides; to investigate fundamental properties of various **few-body systems** and develop **rigorous** and effective **mathematical methods** of calculations of their properties; to explore the dynamics of **particle-nucleus and nucleus-nucleus collisions at relativistic energies** and **nuclear matter properties** and its phase transitions **at extreme values of temperature and density**.

Theory of Condensed Matter

Leaders: V.A. Osipov, A. Povolotskii

Multiparticle models of solids taking into consideration **strong electron correlations, electron-lattice, and spin interactions to describe spectra of quasiparticle excitations, phase transitions and kinetic phenomena in solids.** In **equilibrium and nonequilibrium media** with strong correlations, the processes of **multifragmentation, clusterization in phase transitions and the influence of surface effects** on properties of clusters. In the **theory of superconductivity, nonstandard mechanisms of pairing in metal-oxides,** the problem of bipolaron stability in a polaron gas environment, the influence of **strong electric fields and temperature gradients** on elastic, magnetic, and thermal properties of granular superconductors. **Magnetic ordering in magnetic semiconductors and in metals with a large magnetoresistance.** Theoretical support of the experiments at the **Frank Laboratory of Neutron Physics, JINR.**

Nonlinear problems in multiparticle theory are studied by using modern methods of the renormalization group theory, the **inverse scattering problem, fractal geometry, and the conformal field theory.**

The **microstructure of amorphous state** are studied in the framework of the theoretical models where topological disorder is introduced via **arrays of disclination dipoles and loops.** The thermal properties of disclinated media are of primary interest. The electronic spectrum of **carbon materials, fullerenes and nanotubes,** are examined within the field-theory models adapted to account for nontrivial geometry of these nanostructures. The model of **random Josephson junction arrays** will be studied and applied to describe **high-temperature granular superconductors.**

Modern Mathematical Physics: Strings and Gravity, Supersymmetry, Integrability

A.P. Isaev, A.S. Sorin, S.O. Krivonos, A.T. Filippov

Superstring Theory is the most serious and worldwide pursued candidate for a unified theory of all fundamental interactions including **Quantum Gravity** and thus it is the principal source of the problems which are the subject of modern mathematical physics. The development of the theory involves the study of its surprisingly wide spectrum of possible regimes, vacua and exact classical and quantum solutions. Furthermore, the theory has **applications in many directions including the nonperturbative regime of supersymmetric gauge theories, the mechanics and thermodynamics of black holes and cosmological models of the universe expansion.** These are unique laboratories to check general ideas from unified theories. In particular, in order to accommodate and develop the new ideas in these sectors inspired by String Theory, it is crucial to use the powerful mathematical methods provided by the **theory of Integrable Systems, Quantum Groups and Non-Commutative Geometry.** The goals of the present new theme precisely belong to the bridging between these fields and further development of suitable schemes to be applied in this context.

Dubna International Advanced School of Theoretical Physics (DIAS-TH)

A.S. Sorin, V.V. Voronov

The Bogoliubov Laboratory of Theoretical Physics has a good record of organizing international workshops and schools in Dubna. **DIAS-TH organizes and supervises all educational programs for students, postgraduates, and young scientists at BLTP.** It should function continuously and **the standard short schools (about 3-4 a year) should be organized coherently.** Other educational programs in Dubna such as the **JINR University Center correlate with DIAS-TH** (common programs on modern theoretical physics, workshops for students and young scientists, etc.).

[1996](#) | [1997](#) | [1998](#) | [1999](#) | [2000](#) | [2001](#) | [2002](#) | [2003](#) | [2004](#) | [2005](#) | [2006](#) | [2007](#) | [2008](#) | [2009](#) | [2010](#) | [2011](#) | [2012](#) | [2013](#) | [2014](#) | [2015](#) | [2016](#) | [2017](#) | [2018](#) | [2019](#)

January 28 - February 1
XV Winter School on Theoretical Physics
Complex Systems and Advanced Materials

March 20 - 22
International Workshop
Infinite and finite nuclear matter

June 2 - 8
XIV International Workshop
Particle Correlations and Femtoscopy (WPCF2019)

July 14 - 20
BLTP/JINR - APCTP/Korea Joint Workshop
Modern problems in nuclear and elementary particle physics

July 8 - 12, Prague, Czech Republic
The XXVIth International Conference on
Integrable Systems and Quantum Symmetries

July 22 - August 2
Helmholtz International Summer School
Quantum Field Theory at the Limits: from Strong Fields to Heavy Quarks

July 28 - August 2
International School
Advanced Methods of Modern Theoretical Physics: Integrable and Stochastic Systems

August 4 - 17
Helmholtz International Summer School
19/ smology, Strings, New Physics

August 26 - 31, Yerevan, Armenia
International Workshop
Supersymmetries and Quantum Symmetries (SQS19)

September 1 - 10, Sinaia, Romania
VIIIth International Pontecorvo
Neutrino Physics School

September 2 - 7
XVIIIth International Workshop on
High Energy Spin Physics (DSPIN-19)

September 9 - 13
International Bogolyubov Conference
Problems of Theoretical and Mathematical Physics
(dedicated to the 110th anniversary of the birth of N.N. Bogolyubov (1909-1992))

September 16 - 19
II International workshop
Theory of Hadronic Matter under Extreme Conditions

September 22 – 28, Varna, Bulgaria
XXIII International School on
Nuclear Physics, Neutron Physics and Applications

October 27 - November 1, Guangzhou, China
BLTP/JINR - KLTP/CAS Joint Workshop
Physics of Strong Interacting Systems

Helmholtz International Summer School "COSMOLOGY, STRINGS, NEW PHYSICS"

4-17 August 2019
Europe/Moscow timezone

Overview

Scientific Programme

Registration

Registration Form

Organizers

Travel Info

Visa information

Poster

Support

✉ diastp@theor.jinr.ru

☎ +74962164747

Scientific Programme



Lectures

I.L. Buchbinder (Tomsk Pedagogical Univ.)
"Introduction to supergravity"

D. S. Gorbunov (INR RAS) (3 lectures)
" Introduction to the Hot Big Bang Theory" " Baryogenesis in the Early Universe"
" Dark Matter Models"

D. I. Kazakov (BLTP JINR)
"R-operation and renormalization group for non-renormalizable theories" (2 lectures)

E. T. Musaev (MIPT)
"Branes and string cosmology"

K. A. Postnov (SAI MSU)
"Gravitational waves from black holes and neutron star mergers"

A.A. Starobinsky (Landau Inst. & BLTP JINR)
"Inflation, pre-inflation and post-inflationary heating in modified gravity models" (3 lectures)

V.A. Rubakov (INR RAS) TBA

V.P. Spiridonov (BLTP JINR)
"Superconformal indices, Seiberg dualities and integrable two-dimensional statistical mechanics models " (2 lectures)

A. Vikman (FZU, Prague)
"Dark side of the Universe" (3 lectures)

A. Westphal (DESY)
"Aspects of String Cosmology - de Sitter Vacua and Large-Field Inflation in String Theory" (3-4 lectures)

Conferences and Schools, 2013-2018

Total - 93 (> 5000 participants)

DIAS-TH and Helmholtz Schools - 23

Educational Activities

More than 250 lecture courses at

*JINR University Centre, DIAS-TH, Moscow U.,
Dubna U., MIPhT (SU), etc.*

Research and Education Project DIAS-TH

“Dubna International School of Theoretical Physics”



С 324
Б-74

ОСНОВЫ
АКСИОМАТИЧЕСКОГО
ПОДХОДА
В КВАНТОВОЙ
ТЕОРИИ ПОЛЯ

Н. Н. БОГОЛЮБОВ
А. А. ЛОГУНОВ
И. Т. ТОДОРОВ

Н. Н. БОГОЛЮБОВ, Д. В. ШИРКОВ
ВВЕДЕНИЕ В ТЕОРИЮ
КВАНТОВАННЫХ ПОЛЕЙ

Nikolai Nikolaevich
BOGOLIUBOV

COLLECTION OF SCIENTIFIC WORKS
IN TWELVE VOLUMES

Quantum Theory

Volume 11

N.N. Bogoliubov, A.A. Logunov,
A.I. Oksak, I.T. Todorov

General Principles
of the Quantum Field Theory

MOSCOW NAUKA 2008

Николай Николаевич
БОГОЛЮБОВ

СОБРАНИЕ НАУЧНЫХ ТРУДОВ
В ДВЕНАДЦАТИ ТОМАХ

Квантовая теория

Том XI

Н. Н. Боголюбов, А. А. Логунов,
А. И. Оксак, И. Т. Тодоров

Общие принципы
квантовой теории поля




ФИЗИКА ВЫСОКИХ ЭНЕРГИЙ
И ТЕОРИЯ ЭЛЕМЕНТАРНЫХ ЧАСТИЦ


АЛЛИЛУЕВ С. П.
АНИСОВИЧ В. В.
АНСЕЛЬМ А. А.
АРБУЗОВ Б. А.
АЧАСОВ Н. Н.
БАЛДИН А. М.
БАРБАШОВ Б. М.
БЕРЕСТИЦКИЙ В. Б.
БЛОХИНЦЕВ Д. И.
БОГОЛЮБОВ Н. Н.
ВАЛУЕВ Б. Н.
ВАНЯШИН В. С.
WERLE J.
WICK J.
ВИНТЕРНИЦ П.
ВОЛКОВ Д. В.
VOTRUBA J.
NAVŘÍČEK M.
ГЕЛЬФАНД И. М.
ГЕРШТЕЙН С. С.
ГОВОРКОВ А. Б.
ГРАЕВ М. И.
ДЯТЛОВ И. Т.
ЕФРЕМОВ А. В.
ИОФЕ Б. Л.
ИСАЕВ П. С.
КАДЫШЕВСКИЙ В. Г.
КОБЗАРЕВ И. Ю.
КОМАР А. А.
LIPKIN H. J.
ЛЮМСАДЗЕ Ю. М.
МАКАРОВ А. А.
МАЛЮТА Ю. М.
МАРКОВ М. А.
МАРШАК Р. Е.
МАТВЕЕВ В. А.

МАТВЕЕНКО А. В.
MATTHEWS P. T.
МОЗЖИМАС Я.
МУРАДЯН Р. М.
НДИМАРК М. А.
НГУЕН ВАН ХЪЕУ
НОВОЖИЛОВ Ю. В.
ОГНЕВЕЦКИЙ В. И.
ОКОНОВ Э. О.
OLSZEWSKI J.
ПИСАРЕНКО В. Г.
ПОДВИВАНОВ М. К.
ПОДУБАРИНОВ В. И.
ПОПОВ В. С.
ROSENFELD A.
СОЛОВЬЕВ Л. Д.
СТАВРАКИ Г. Л.
СТОЯНОВ Д. П.
СТРУМИНСКИЙ Б. В.
ТАВХЕЛИДЗЕ А. И.
ТЕРЕНТЬЕВ И. А.
ТОДОРОВ И. Т.
UHELMANN A.
ФАДДЕЕВ Л. Д.
ФИЛИППОВ А. Т.
FRONSDAL S.
ЧЕРНИКОВ Н. А.
ШЕЛЕСТ В. П.
ШИРКОВ Д. В.



МОСКВА НАУКА





V. Inozemtsev
Integrable Heisenberg-Dirac chains ...

V. Pervushin, A. Pavlov,
Principles of Quantum Universe

Madimir Inozemtsev
Integrable Heisenberg-Dirac chains with variable range exchange
 integrable quantum spin chains



A.P. Isaev and V.A. Rubakov,
Theory of Groups and Symmetries. Finite Groups, Lie Groups and Lie Algebras.

Theory of Groups and Symmetries
 Finite Groups, Lie Groups, and Lie Algebras



Theory of Groups and Symmetries



The book presents the main approaches in study of algebraic structures of symmetries in models of theoretical and mathematical physics, namely groups and Lie algebras and their deformations. It covers the commonly encountered quantum groups (including Yangians). The second main goal of the book is to present a differential geometry of coset spaces that is actively used in investigations of models of quantum field theory, gravity and statistical physics. The third goal is to explain the main ideas about the theory of conformal symmetries, which is the basis of the AdS/CFT correspondence.

The theory of groups and symmetries is an important part of theoretical physics. In elementary particle physics, cosmology and related fields, the key role is played by Lie groups and algebras corresponding to continuous symmetries. For example, relativistic physics is based on the Lorentz and Poincare groups, and the modern theory of elementary particles — the Standard Model — is based on gauge (local) symmetry with the gauge group $SU(3) \times SU(2) \times U(1)$. This book presents constructions and results of a general nature, along with numerous concrete examples that have direct applications in modern theoretical and mathematical physics.

Isaev
 Rubakov

Alexey P Isaev
 Valery A Rubakov



Виктор Пераушин
 Александр Павлов

Принципы квантовой Вселенной



World Scientific
 worldscientific.com



World Scientific

Принципы квантовой Вселенной


Пераушин, Павлов

AGREEMENTS

- **BLTP – ICTP** (since '88)
- **BLTP – Germany** (since '91)
Heisenberg-Landau Program
- **BLTP – INFN** (since XII '95)
6 month visits to Italy
- **BLTP – CERN-TH** (since XII '95)
3 month visits to CERN
- **BLTP – Poland** (since XII '98)
Bogoliubov-Infeld Program
- **BLTP – Czech Republic** (since XII '99)
Blokhintsev-Votruba Program
- **BLTP – Romania** (since XII '03)
Titeica-Markov Program
- **BLTP – APCTP, Pohang** (since '07)
- **BLTP – Bulgaria** (since '09)
Soloviev-Khristov Program
- **BLTP – ITP CAS, China** (since VII '10)
- **BLTP – IOP VAST, Vietnam** (since VIII '11)
- **BLTP – Physical Inst., NAS, Armenia** (since '09)
Smorodinsky - Ter-Antonyan Program
- **BLTP- IMSc India** (since 2016)

THEORY OF FUNDAMENTAL INTERACTIONS

The state of art analysis of the Stability of the Standard Model vacuum



Based on record 3-loop order calculations it was shown that for a fixed value of $M_h=125.7$ GeV absolute SM stability leads to a bound on the top mass

$$M_t < M_t^{crit} = 171.54 \text{ GeV}$$


compatible with measured $M_t^{expt} = 173.21(87)$ GeV at the 1.3 sigma level. Previously made conclusion that the SM vacuum is metastable may be premature.

A. Bednyakov et al, *Phys. Rev. Lett.* 115, 201802 (2015);
JHEP 1301 (2013) ; *Phys.Lett.* B722 (2013); *Nucl. Phys.* B875 (2013).

Constraints on Supersymmetry from LHC data combined with cosmology and direct dark matter searches

A combined LHC data (LHCb experiment), the relic density (WMAP and other cosmological data) and upper limits on the dark matter scattering cross sections on nuclei (XENON100 data) indicate that gluinos below 1 TeV and the lightest supersymmetric particle with the mass below 160 GeV are excluded.


C. Beskidt, W. de Boer, D.I. Kazakov, F. Ratnikov, *Eur. Phys. J. C* (2012); *JHEP* (2012)



BaBar puzzle

BaBar puzzle for pion transition form factor was systematically and comprehensively analyzed using the various QCD-motivated approaches. It is concluded that the current experimental accuracy (BaBar and Belle experiments) is not sufficient to discriminate between QCD factorization and its violation.

A.P. Bakulev, S.V. Mikhailov, A.V. Pimikov, N.G. Stefanis, *Phys. Rev. D* 86 (2012); *Few Body Syst.* 55 (2014); **A. Dorokhov, E.Kuraev**, *Phys. Rev.D* 88(2013); Ya.Klopot, A.Oganesian, O.Teryaev, *Phys.Rev.D* 87 (2013).



Determination of the electron-to-proton mass ratio


The record 10-fold improvement for the transition energies of the hydrogen molecular ion and the antiprotonic helium was obtained. This allows one to achieve a fractional precision of $1.5 \cdot 10^{-11}$ in determination of the electron-to-proton mass ratio. This progress may have serious impact on the fundamental physical constants.

V.Korobov, et al, *Phys.Rev.Lett.* 112 (2014).

Explanation of the “reactor anomaly”

Within the field theoretical approaches to neutrino flavor oscillations It is shown that for relatively short but macroscopic baselines, there must be a small violation of the classical inverse-square law for the neutrino flux. Numerical results provide an explanation of the long-standing “reactor anomaly”.

V.A. Naumov, D.S. Shkirmanov, *Mod. Phys. Lett. A30* (2015); *Eur. Phys. J. C73* (2013).




HADRONIC MATTER UNDER EXTREME CONDITIONS

Helicity separation in heavy-ion collisions


The P-odd effects related to the vorticity of the medium formed in noncentral heavy-ion collisions are studied. The noticeable hydrodynamical helicity was observed to manifest specific mirror behavior with respect to the reaction plane. The effect is maximal at the NICA and FAIR energy range.

M. Baznat, K. Gudima , A. Sorin, O. Teryaev, *Phys.Rev. C88* (2013).



Polarization of the QCD vacuum by the strong electromagnetic fields

The quark interaction with both confining vacuum gluon fields and external strong electromagnetic field created in HIC leads to specific polarization of the vacuum gluon fields, and EM field may trigger deconfinement in HIC. This polarization effect was confirmed later in Lattice QCD calculations (LQCD - *JHEP* 1304 (2013); *Phys. Rev. Lett.* 110 (2013)).




B. Galilo, S. Nedelko, *Phys.Rev.D* 84 (2011) ; S. Nedelko, V.Voronin, *Eur.Phys. J. A* (2015).

Finite temperature gluon spectral functions from $N_f=2+1+1$ lattice QCD

For the first time gluon spectral functions at finite temperature in Landau gauge were calculated based on lattice QCD with $N_f=2+1+1$ dynamical quark flavors combined with a novel Bayesian approach for the extraction of non-positive definite spectral functions. Gluon quasi-particle peaks in the spectral function were found. The obtained gluon dispersion relation reveals different in-medium masses for longitudinal and transversal gluons. This methodological advances allows one to perform ab initio QCD studying of the transport properties of quark-gluon matter, which is highly important for theoretical description of the relativistic heavy ion physics.


E.-M. Ilgenfritz, J. M. Pawlowski, A. Rothkopf, A. Trunin, *ArXiv:1701.08610[hep-lat]*



Proton spin calculation within Lattice QCD

For the first time the contribution of orbital angular momenta of up, down and strange quarks and gluon angular momenta to proton spin were computed using LQCD. It has been found that they contribute ~47% (sum of individual quark contribution) and ~28%, respectively, and thus accounting for all of missing proton spin. The other quantity, quark spin contribution, has been known for a long time to be ~25% both experimentally and in Lattice QCD.

M. Deka et al [χ QCD collaboration] *Phys. Rev. D* 91 (2015)





THEORY OF NUCLEAR STRUCTURE AND NUCLEAR REACTIONS

β -decay half-lives of the neutron-rich $N = 50$ isotones

It was shown that tensor interactions and 2p-2h configuration result in decreasing of beta-decay half lives of the neutron-rich $N = 50$ isotones significantly. Results of the calculations without the tensor interaction (open triangles, circles, squares) and **with the tensor interaction (filled triangles, circles, squares)** are shown.

Experimental data - filled diamonds.

Severiukhin A., Voronov V., Borzov I., Arseniev N., Nguen Van Giai, *Phys. Rev. C90* (2014).



Nuclear weak interaction processes in supernova


A new method to compute rates and cross sections for nuclear weak interaction processes in supernova environments was developed. For iron isotopes the results demonstrate that temperature shifts the Gamow-Teller (GT) strength function to lower energies, thereby leading to a significantly larger rate increase than expected from the shell model calculations. The method was also applied to study supernova neutrino nucleus reactions.

A.A. Dzhiyev, et al., *Phys. Rev. C81* (2010); *Phys. Rev. C89* (2014); *Phys. Rev. C92*(2015)


Cluster approach for describing nuclear fission

The model is developed suited for the description of the symmetric and asymmetric charge yields, as well as the transition between them. Unexpectedly large asymmetric mode for the fission of Th and Ra isotopes was observed at high excitation energy. Triple-peak charge distributions are predicted for fissioning $N \sim 136$ nuclei.

Pasca H., Andreev A.V., Adamian G.G., Antonenko N.V., *PRC* 93, 034620 and 054602 (2016); 94, 064614 (2016); *PLB* 760, 800 (2016); *EPJA* 52, 369 (2016)



Predictions of shell closures in the region of unknown nuclei




It is shown that the two-center shell model indicates the magic nucleus next to ^{208}Pb has the charge $Z \geq 120$ and the half-life $1,7\text{ms} - 0,16\text{s}$ (for $Z=120, N=175-179$)

Calculated (symbols connected by lines) and experimental energies of α decay of the nuclides with even Z from the range $108 \leq Z \leq 126$ are shown in figure.

G.G.Adamian, N.V.Antonenko, R.V.Jolos, A.K.Nasirov, V.G.Kartavenko, L.A.Malov, T.M.Shneidman, A.Sushkov, A.N.Bezbakh, *Eur. Phys. J. A45 (2010); Eur. Phys. J. A47 (2011); Phys. Rev. C 84 (2011); Phys. Rev. C 85 (2012); Phys. Rev. C 91 (2015).*

Discovery of confinement-induced resonances




A splitting of the confinement-induced resonance in ultracold gases was predicted and then discovered in experiment. The developed theory has allowed to describe quantitatively the resonance effects: width and shifts of the s-wave CIR in an atomic trap and its splitting caused by the trap anisotropy. Predictions of the theory – the widths and shifts of the p-wave CIRs and the found conditions for appearance of dipole CIRs – are extremely relevant to the ongoing research of cold atoms and molecules.

V.S. Melezhik, et al, *Phys. Rev. Lett. 104 (2010); Phys. Rev. A 84, 042712 (2011); Phys. Rev. A 86, 062713 (2012); Phys. Rev. Lett. 111, 183201 (2013); J. Phys. B 48, 155301 (2015).*

The electron-positron pair emission in interaction of the photon with laser pulses

For the first time it has been found that the short laser pulses “generate” high momentum components which produce a great amplifier effect for the multi-photon, sub-threshold events. Enhancement may reach many orders of magnitude depending on field intensity and the beam shape. The probability of the electron-positron pair emission in interaction of the probe photon with the short and intensive laser pulses is enhanced (by several orders of magnitudes).

A. Titov, et al, *Phys.Rev. A 87 (2013); Phys. Rev. Lett. 108, (2012); Eur. Phys. J. D 68 (2014).*



CONDENSED MATTER THEORY

Graphene tunnel field-effect transistor

A new concept for a graphene tunnel field-effect transistor was proposed. The main idea is based on the use of two graphene electrodes with zigzag termination divided by a narrow gap under the influence of the common gate. It was shown that the device will have a pronounced switching effect at low gate voltage and high on/off current ratio at room temperature.


V. Osipov, V. Katkov, *Applied Phys. Lett.* 104 (2014) 053102; *JETP Lett.* 98 (2013) 782; *Patent for invention No 2554694 [01.06.2015].*



Surface fractals and the small-angle neutron scattering

New general properties of fractals have been predicted. Particularly important result is that a surface fractal can be realized as a sum of mass fractals, and thus the small-angle neutron scattering amplitude of the surface fractal can be calculated as a sum of the mass fractal amplitudes.

E. M. Anitas *et al*, *Rom. Journ. Phys.* 60, 658 (2015); *J. Appl. Cryst.* 47, 198 (2014); *Phys. Rev. E* 84, 036203 (2011).



Microscopic theory of metals with strong electron correlations

A microscopic theory of metals with strong electron correlations was developed. The theory is based on a consistent consideration of kinematic interaction of electrons with spin and charge fluctuations. Within this approach, a microscopic description of high temperature superconductivity in cuprates without fitting coupling constant, a theory of magnetic excitations in the normal and superconducting states, and a theory of electrical conductivity and optical absorption have been created.

N. M. Plakida, *High-Temperature Cuprate Superconductors*, Springer, 2010; N. M. Plakida, V. S. Oudovenko, *Eur. Phys. J. B* 86 (2013); N. M. Plakida, V. S. Oudovenko, *JETP* 146 (2014); A. A. Vladimirov, D. Ihle, N. M. Plakida, *Phys. Rev. B* 83 (2011); A. A. Vladimirov, D. Ihle, N. M. Plakida, *Phys. Rev. B* 85 (2012).

Emergence of jams in ASEP


An integrable generalization of totally asymmetric simple exclusion process (ASEP) with extra interaction was constructed. The distribution of particle current shows two types of universal scaling behaviour depending on the value of the interaction strength. Either, when the density of particle clusters is finite, the distribution has the form specific for Kardar–Parisi–Zhang universality class, or all particles form a single cluster, characterized by the Gauss distribution. Unification of the two universal regimes and description of the transition between them were obtained.

A.E. Derbyshev, A.M. Povolotsky, V.B. Priezzhev, *Phys. Rev. E* 91 (2015).

Strongly frustrated triangular spin lattice emerging from triplet dimer formation in honeycomb Li_2IrO_3


Iridium oxides with a honeycomb lattice have been identified as platforms for the much anticipated Kitaev topological spin liquid. A new type of magnetic ground state that was called “the structure of triplet dimers on an effective triangle lattice” has been predicted. The prediction can be verified by means of neutron diffraction and neutron magnetic spectroscopy.

S. Nishimoto, V. Katukuri, V. Yushankhai, H. Stoll, U. Roessler, L. Hozoi, I. Rousochatzakis, and J. van den Brink, *NATURE Comm.* 7, 10273 (2016)



MODERN MATHEMATICAL PHYSICS

HYM-flation: Yang–Mills cosmology with Horndeski coupling



It is discovered that the effective gravity theory with non-minimal (Horndeski) coupling of gauge fields to gravity gives rise to robust Planck-scale inflation with graceful exit. This solves the initial conditions problem for the observed GUT-scale inflation. Considering both gauge and Higgs fields one obtains a consistent description of the early Universe based solely on the fields of Standard Model.

E. Davydov, D. Gal'tsov, *Phys. Lett. B* 753, 622 (2016)

New proof of the Atiyah-Singer theorem

The full description of the geometries of the most general $N=2$ supersymmetric quantum mechanical models was done and a new simple physical proof of the Atiyah-Singer theorem was given. A new example of $N=2$ supersymmetric quantum mechanical sigma models was studied. A special class of quasicomplex sigma models was derived. These results are important for understanding of the properties of quantum states in the field theories with anomalies and supersymmetric theories in diverse dimensions.

S. Fedoruk, E. Ivanov, A. Smilga, *Int. J. Mod. Phys. A* 27(2012)1230024, A27(2012)1250146; *SIGMA* 9(2013)069; *J. Math. Phys.* 55(2014)052302; *J. Phys.* A48(2015)215401.

N=4 supersymmetric mechanics on curved spaces

We present $N = 4$ supersymmetric mechanics on n -dimensional Riemannian manifolds constructed within the Hamiltonian approach. The structure functions entering the supercharges and the Hamiltonian obey modified covariant constancy equations as well as modified Witten–Dijkgraaf–Verlinde–Verlinde equations specified by the presence of the manifold's curvature tensor. Solutions of original Witten–Dijkgraaf–Verlinde–Verlinde equations and related prepotentials defining $N=4$ superconformal mechanics in flat space can be lifted to $so(n)$ -invariant Riemannian manifolds.

N.Kozyrev, S.Krivonos, O.Lichtenfeld, A.Nersessian, A. Sutulin,
Phys.Rev. D97 (1918) 085015

Elliptic hypergeometry of supersymmetric dualities

Seiberg type dualities were constructed for supersymmetric field theories outside the conformal window. Explicit forms of the superconformal indices were derived in terms of the elliptic hypergeometric integrals and systematically studied for a number of supersymmetric theories.

V.P. Spiridonov, G.S. Vartanov, *Phys. Rev. Lett.* 105 (2010), *Comm. Math. Phys.* 304 (2011); *Comm. Math. Phys.* 325 (2014); *JHEP* 06 (2012) 016; *JHEP* 06 (2014), 062; *Contemp. Math.* 563 (2012), 181-211.





DUBNA

JINR

BLTP

Welcome!





| [1996](#) | [1997](#) | [1998](#) | [1999](#) | [2000](#) | [2001](#) | [2002](#) | [2003](#) | [2004](#) | [2005](#) | [2006](#) | [2007](#) |
[2008](#) | [2009](#) | [2010](#) | [2011](#) | [2012](#) | [2013](#) | [2014](#) | [2015](#) | [2016](#) | [2017](#) | [2018](#)

January 29 - February 2

XIVth Winter School on Theoretical Physics
Partition Functions and Automorphic Forms

April 16 - 18

II International Workshop
Simulations of HIC for NICA energies

April 18 - 19

Colloquium on
Nonequilibrium phenomena in strongly correlated systems

May 27 - June 2, Roshchino, Valdai District, Russia
20th International Seminar on
High Energy Physics "Quarks-2018"

June 3 - 9, Bourgas, Bulgaria

International Conference
Nuclear Structure and Related Topics (NSRT18)

July 9 - 12

International workshop
Low Dimensional Materials: Theory, Modelling and Experiment

July 9 - 13, Prague, Czech Republic

The 32nd International Colloquium on
Group Theoretical Methods in Physics

July 22 – August 1

Helmholtz International Summer School
Modern Colliders - Theory and Experiment 2018
and Workshop
Calculations for Modern and Future Colliders

August 5 - 10

International School Advanced Methods of Modern Theoretical Physics:
Integrable and Stochastic Systems

August 13 - 16

International Workshop
Supersymmetry in Integrable Systems (SIS'18)

August 20 - 31

Helmholtz International Summer School
Matter under Extreme Conditions in Heavy-Ion Collisions and Astrophysics

August 27 - 31, Peterhof, Russia

VI International Conference
Models in Quantum Field Theory (MQFT-2018)
in memory of prof. A.N. Vasiliev

September 2 - 7, Saint-Petersburg, Russia

BLTP/JINR - KLFTP/CAS Joint Workshop
Physics of Strong Interacting Systems

September 4 - 6

II International Workshop
Lattice and Functional Techniques for Exploration of Phase Structure and Transport Properties in Quantum Chromodynamics

September 24 - 27, INR RAS, Moscow, Russia

XV International Seminar on
Electromagnetic Interactions of Nuclei (EMIN-2018)

September 17 - 22

The XXIV International Baldin Seminar on High Energy Physics Problems
Relativistic Nuclear Physics and Quantum Chromodynamics